

CLINICAL AND EXPERIMENTAL OBSTETRICS & GYNECOLOGY

an International Journal

Founding Editor

A. Onnis

Montréal (CND)

Editors-in-Chief

M. Marchetti

Montréal (CND)

J.H. Check

Camden, NJ (USA)

Assistant Editor

J. Wilson

San Diego - CA (USA)

Editorial Board

Audet-Lapointe P., *Montréal (Canada)*

Axt-Fliedner R., *Lübeck (Germany)*

Basta A., *Krakow (Poland)*

Bender H.J., *Dusseldorf (Germany)*

Bhattacharya N., *Calcutta (India)*

Bonilla Musoles F., *Valencia (Spain)*

Charkviani T., *Tbilisi (Georgia)*

Dexeus S., *Barcelona (Spain)*

Di Paola G., *Buenos Aires (Argentina)*

Eskes T.K.A.B.,

Nijmegen (The Netherlands)

Farghaly S.A., *New York (USA)*

Friedrich M., *Homburg (Germany)*

Gomel V., *Vancouver (Canada)*

Gorins A., *Paris (France)*

Grella P.V., *Padua (Italy)*

Holub Z., *Kladno (Czech Republic)*

Jordan J.A., *Birmingham, England (UK)*

Kaplan B., *Petach Tikva (Israel)*

Kralj B., *Ljubljana (Slovenia)*

Markowska J., *Poznan (Poland)*

Marth C., *Innsbruck (Austria)*

Meden-Vrtovec H., *Ljubljana (Slovenia)*

Ohara N., *Kobe (Japan)*

Papadopoulos N., *Alexandroupolis (Greece)*

Rakar S., *Ljubljana (Slovenia)*

Sciarra J.J., *Chicago, IL (USA)*

Stelmachow J., *Warsaw (Poland)*

Varras M.N., *Athens (Greece)*

Winter R., *Graz (Austria)*

Publishing Organization (M. Morsani):

I.R.O.G. CANADA, Inc. - 4900 Côte St-Luc - Apt # 212 - Montréal, Qué. H3W 2H3 (Canada)

Tel. +514-4893242 - Fax +514-4854513 - E-mail: canlux@mgroun-online.com - www.irog.net

Editorial Office (M. Critelli):

Galleria Storione, 2/A - 35123 Padua (Italy) - Tel. (39) 049 8756900 - Fax (39) 049 8752018

CLINICAL AND EXPERIMENTAL OBSTETRICS AND GYNECOLOGY (ISSN 0390-6663) publishes original work, preferably brief reports, in the fields of Gynecology, Obstetrics, Fetal Medicine, Gynecological Endocrinology and related subjects. (Fertility and Sterility, Menopause, Uro-gynecology, Ultrasound in Obstetrics and Gynecology, Sexually Transmitted Diseases, Reproductive Biological Section). The Journal is covered by **INDEX MEDICUS, MEDLINE, EMBASE/Excerpta Medica**.

CLINICAL AND EXPERIMENTAL OBSTETRICS AND GYNECOLOGY is issued every three months in one volume per year by IROG CANADA Inc. Montréal. Printed in Italy by "La Garangola", Tipografia Editrice - Via E. Dalla Costa, 6 - 35129 Padova (Italy).

EDITORIAL ARTICLES

- A practical approach to the prevention of miscarriage: Part 5 – antiphospholipid syndrome as a cause of spontaneous abortion** 5
 J.H. Check - *Camden, NJ (USA)*
 Antiphospholipid syndrome is a remediable (with anticoagulants) cause of pregnancy loss, especially spontaneous abortion after ten weeks.
- Chronic pelvic pain syndromes – traditional and novel therapies: Part I surgical therapy** 10
 J.H. Check - *Camden, NJ (USA)*
 Laparoscopic ablation of endometriosis, surgical excision of endometriosis, laparoscopic uterosacral nerve ablation, presacral neurectomy, and total abdominal hysterectomy with bilateral salpingectomy have a secondary role in treating pelvic pain.

REVIEW ARTICLE

- Maternal obesity and pregnancy outcome** 14
 K. Sameera Begum, K. Sachchithanantham, S. Somsubhra De - *Melaka, MALAYSIA*
 The increasing prevalence of obesity in young women is a major public health concern. The trends have a major impact not only on pregnancy outcome but also later in life to both mother and children. Education about achieving a healthy body weight before conception is important.

ORIGINAL ARTICLES

Reproductive Biology Section

- About 13% of women may have the wrong method of oocyte insemination when undergoing in vitro fertilization by failure to evaluate the abnormal hypo-osmotic swelling test score** 21
 G. Citrino, J.H. Check, A. Bollendorf, W. Hourani, A. Diantonio - *Camden, NJ (USA)*
 A retrospective study found that 12.5% of the male partners of couples undergoing in vitro fertilization have low hypo-osmotic swelling test scores despite otherwise normal semen parameters.
- Comparison of the efficacy of treating sperm with low hypoosmotic swelling test scores with chymotrypsin followed by intrauterine insemination vs in vitro fertilization with intracytoplasmic sperm injection** 24
 A. Bollendorf, D. Check, J.H. Check, W. Hourani, K. McMonagle - *Camden, NJ (USA)*
 One cycle of in vitro fertilization with intracytoplasmic sperm injection is much more effective than the treatment of sperm with chymotrypsin followed by two cycles of intrauterine insemination.
- Comparison of pregnancy outcome following frozen embryo transfer (ET) in a gestational carrier program according to source of the oocytes** 26
 J.H. Check, B. Katsoff, D. Brasile, C. Wilson, D. Summers-Chase - *Camden, NJ (USA)*
 A retrospective study found good pregnancy outcome following frozen embryo transfer into gestation carriers whether the source of oocytes was from fertile egg donors or the infertile female partner.

General Section

- Chronic action of association of zidovudine, lamivudine and ritonavir on pregnant rats. A biologic assay** 28
 A. Wagner, M. Uchiyama Nakamura, R.S. Simões, R.M. Oliveira-Filho, T.M. Pereira Fontes, L.P. Fogarolli de Carvalho, S. Espiridiao, L. Kulay Jr. - *São Paulo, SP, BRAZIL*
 The administration of zidovudine-lamivudine-ritonavir in rat pregnancy can reduce the maternal body weight gain and lower the term placental weight.

| | |
|--|----|
| A comparative study between microwave endometrial ablation and conventional surgical procedures for treatment of menorrhagia | 33 |
| K. Nakayama, S. Yeasmin, A. Katagiri, M.T. Rahman, M. Rahman, M. Ishikawa, K. Iida, N. Nakayama, S. Aoki, K. Miyazaki - <i>Shimane, JAPAN</i> MEA is a promising new method for treating menorrhagia instead of conventional surgical procedures. | |
| Apoptosis and expression of Bcl-2, Bax, p53, caspase-3, and Fas, Fas ligand in placentas complicated by preeclampsia | 38 |
| I. Mendilcioglu, S. Karaveli, G. Erdogan, M. Simsek, O. Taskin - <i>Antalya, TURKEY</i> Increased apoptosis in trophoblasts may not be associated with significant alterations in regulator proteins. | |
| Doppler assessment between pathological examination of the placenta and late fetal intrauterine demise | 43 |
| I. Babovic, J. Tadic, S. Plesinac, Z. Radojicic, D. Plecas - <i>Belgrade, SERBIA</i> Umbilical artery Doppler is a poor predictor of stillbirths due to placental dysfunction. | |
| Intravaginal misoprostol reduces intraoperative blood loss in minimally invasive myomectomy: a randomized clinical trial | 46 |
| I. Kalogiannidis, P. Xiromeritis, N. Prapas, Y. Prapas - <i>Thessaloniki, GREECE</i> Preoperative use of misoprostol reduces intraoperative blood loss in patients managed by minimally invasive myomectomy. | |
| The golden ratio of nasal width to nasal bone length | 50 |
| G. Goynumer, M. Yayla, B. Durukan, L. Wetherilt - <i>Istanbul, TURKEY</i> Calculating the ratio of fetal nasal width over nasal bone length might be useful in evaluating fetal abnormalities. | |
| The effect of low-dose combined oral contraceptive containing 100 ug levonorgestrel on plasma plasminogen activator inhibitor-1 concentrations | 54 |
| M. Küçük, S.D. Sezer, A.R. Odabaşı, Z. Güner, H. Yüksel, M. Serter - <i>Aydin, TURKEY</i> Similar plasma PAI-1 concentrations were observed in the levonorgestrel/ethinyl estradiol group compared with the control group. | |
| A prospective randomized study for evaluation of wound retractors in the prevention of incision site infections after cesarean section | 57 |
| T.D. Theodoridis, K.N. Chatzigeorgiou, L. Zepiridis, A. Papanicolaou, D. Vavilis, F. Tzevelekis, B.C. Tarlatzis - <i>Thessaloniki, GREECE</i> Results of a study on the use of a new retractor for prevention of wound infections during cesarean section are reported. | |
| Troponin I, C-reactive protein and fibrinogen levels in missed abortions | 60 |
| L. Mollamahmutoğlu, Ş. Kalyoncu, Y. Engin-Üstün, Ö. Moraloğlu, R. Deveer, N. Danışman, U. Buyukcaginci - <i>Ankara, TURKEY</i> C-reactive protein levels are increased in women with missed abortions. | |
| Emergent surgical treatment of radiation-induced enteropathies for patients with urogynecological and colorectal carcinomas | 63 |
| A. Parlakgumus, K. Caliskan, H. Ayse Parlakgumus, F. Kayaselcuk, A. Ezer, T. Colakoglu, S. Belli, S. Yildirim - <i>Ankara, TURKEY</i> Ten years of experience of emergent surgical treatment of radiation enteritis for patients with urogynecological and colorectal carcinomas is reported. | |
| Ectopic pregnancy; risk factors and comparison of intervention success rates in tubal ectopic pregnancy | 67 |
| M. Kazandi, V. Turan - <i>Izmir, TURKEY</i> Ectopic pregnancy is one of the most common diseases in gynecology; its risk factors and management should be well known. | |
| Multiparity, perinatal morbidity and mortality | 71 |
| A. Andrejevic, S. Cvetkovic, Z. Vitosevic, L. Andrejevic, G. Relic - <i>Krusevac, SERBIA</i> Multiparity, perinatal morbidity and mortality in an Albanian population were studied. | |
| Effect of GnRH antagonist therapy on the expression of MUC-1 and heparin binding growth factor expression in the endometrium of hyperstimulated rats | 76 |
| H.T. Ozcakir, A.G. Taman, C. Kose, K. Ozbilgin, S. Inan, H. Caglar - <i>Izmir, TURKEY</i> Results of the study suggest that GnRH antagonist influence the expression of heparin binding growth factor and MUC-1 in the endometrium. | |

| | |
|---|-----|
| Low molecular weight heparin and first trimester maternal PAPP-A and hCG levels, fetal nuchal translucency in the first trimester of pregnancy | 81 |
| E. Tarim, T. Cok, S. Hacivelioglu, T. Bagis - <i>Ankara, TURKEY</i> | |
| Women who used LMWHs during their pregnancy have decreased nuchal translucency compared with unaffected women. | |
| CASE REPORTS | |
| A case of disseminated peritoneal leiomyomatosis and diffuse uterine leiomyomatosis | 84 |
| G. Rosica, G. Santilli, D. Bucari, B. Amici, F. Bulletti, F. Patacchiola, A. Spagnoli, G. Falcocchio - <i>Rieti, ITALY</i> | |
| The diagnosis, differential diagnosis and management of a rare case of disseminated peritoneal leiomyomatosis and diffuse uterine leiomyomatosis are reported. | |
| Marked improvement of headaches and vasomotor symptoms with sympathomimetic amines in a woman with sympathetic hyperalgesia-edema syndrome | 88 |
| J.H. Check, R. Cohen - <i>Camden, NJ (USA)</i> | |
| The beneficial effect of sympathomimetic amines on migraines may not necessarily be related to improving edema. | |
| Lost intrauterine contraceptive device inserted 42 years before: a case report | 90 |
| S.D. Sezer, A.R. Odabaşı, M. Küçük, H. Yüksel - <i>Aydin, TURKEY</i> | |
| In a asymptomatic patient with an IUD localized in the abdomen, the risks of both operating and not operating should be discussed thoroughly. | |
| Labial fusion first diagnosed during pregnancy with voiding difficulty and its management: a case report | 94 |
| M. Küçük, S. Halil, F. Ocer, E. Oral - <i>Istanbul, TURKEY</i> | |
| The case of a 24-year-old woman with extensively fused labia and in the 6th week of pregnancy is reported together with the management. | |
| Placement of a vena cava filter in term pregnancy: case report and review of the literature | 96 |
| B. Zeybek, M.C. Terek, C. Guven, C. Cinar - <i>Izmir, TURKEY</i> | |
| Placement of an inferior vena cava filter in a term pregnant woman with deep venous thrombosis. | |
| Successful management of a high-risk pregnancy with polyhydramnios, IUGR and recurrent pregnancy loss in a chronic renal failure patient: a case report | 99 |
| C. Baykal, S. Kaya, M.K. Takal, Ü. Yakupoğlu - <i>Istanbul, TURKEY</i> | |
| A case of the successful management of a high-risk pregnancy with polyhydramnios, IUGR and with a history of recurrent pregnancy loss in a chronic hemodialysis patient is described. | |
| Infected tuboovarian hydatid cyst: a rare cause of tuboovarian abscess | 102 |
| H. Ayse Parlakgumus, A. Parlakgumus, B. Haydardedeoğlu, F. Bolat - <i>Ankara, TURKEY</i> | |
| A rare cause of tuboovarian abscess in a patient with an infected tuboovarian hydatid cyst is reported. | |