

europaean journal of gynaecological oncology

an International Journal

Editors-in-Chief

A. Onnis **P. Bószé**
Montréal (Canada) Budapest (Hungary)

Associate Editor

M. Marchetti
Montréal (Canada)

Assistant Editor

J. Wilson
San Diego - CA (USA)

Editorial Board

Allen H.H., <i>London, Ontario (Canada)</i>	Geisler H.E., <i>Indianapolis, IN (USA)</i>	Monsonego J., <i>Paris (France)</i>
Anteby S.O., <i>Jerusalem (Israel)</i>	Gerbault A., <i>Villejuif (France)</i>	Pálfalvi L., <i>Budapest, (Hungary)</i>
Audet-Lapointe P., <i>Montreal (Canada)</i>	Gorins A., <i>Paris (France)</i>	Pinotti A., <i>São Paulo (Brazil)</i>
Ayhan A., <i>Ankara (Turkey)</i>	Heintz A.P.M., <i>Utrecht (The Netherlands)</i>	Piura B., <i>Beer Sheva (Israel)</i>
Balat O., <i>Graziantep (Turkey)</i>	Ioannidou-Mouzaka L., <i>Athens (Greece)</i>	Piver S.M., <i>Buffalo, NY (USA)</i>
Barber H.K., <i>New York, NY (USA)</i>	Jordan J.A., <i>Birmingham, England (UK)</i>	Rakar S., <i>Ljubljana (Slovenia)</i>
Basta A., <i>Krakow (Poland)</i>	Karseladze A.I., <i>Moscow (Russia)</i>	Raju K.S., <i>London, England (UK)</i>
Bender H.C., <i>Dusseldorf (Germany)</i>	Klastersky J., <i>Bruxelles (Belgium)</i>	Shepherd J.H., <i>London, England (UK)</i>
Benedetti Panici P.L., <i>Rome (Italy)</i>	Kubista E., <i>Vienna (Austria)</i>	Stelmachów J., <i>Warsaw (Poland)</i>
Charkviani T., <i>Tbilisi (Georgia)</i>	Kudelka A.P., <i>Houston, TX (USA)</i>	Tjalma W., <i>Antwerpen (Belgium)</i>
De Oliveira C.F., <i>Coimbra (Portugal)</i>	Maggino T., <i>Padua (Italy)</i>	Ungár L., <i>Budapest (Hungary)</i>
Dexeus S. Jr., <i>Barcelona (Spain)</i>	Markowska J., <i>Poznan (Poland)</i>	Vermorken J.B., <i>Edegem (Belgium)</i>
Di Paola G.R., <i>Buenos Aires (Argentina)</i>	Marth C., <i>Innsbruck (Austria)</i>	Vírtej P., <i>Bucharest (Romania)</i>
Di Re F., <i>Milan (Italy)</i>	Massuger Leon F.A.G., <i>Nijmegen (The Netherlands)</i>	Wang P.-H., <i>Taipei (Taiwan)</i>
Di Saia P., <i>Orange, CA (USA)</i>	Menczer J., <i>Savyon (Israel)</i>	Winter R., <i>Graz (Austria)</i>
Friedrich M., <i>Hamburg (Germany)</i>		Yokoyama Y., <i>Hirosaki (Japan)</i>
Fuller A.F. Jr., <i>Boston, MA (USA)</i>		

Publishing Organization (M. Morsani):

I.R.O.G. CANADA, Inc. - 4900 Côte St-Luc - Apt # 212 - Montréal, Qué. H3W 2H3 (Canada)
Tel. +514-4893242 - Fax +514-4854513 - E-mail: canlux@mgroun-online.com - www.irog.net

Editorial Office (M. Critelli):

Galleria Storione, 2/A - 35123 Padua (Italy) - Tel. (39) 049 8756900 - Fax (39) 049 8752018

EUROPEAN JOURNAL OF GYNAECOLOGICAL ONCOLOGY (ISSN 0392-2936) publishes original peer reviewed works in the fields of female genital cancers and related subjects and also proceedings of gynecologic oncology society meetings all over the world. The Journal is covered by **CURRENT CONTENTS, SCISEARCH, RESEARCH ALERT, INDEX MEDICUS, MEDLINE, EMBASE/Excerpta Medica, CURRENT ADVANCES IN CANCER RESEARCH, BIOSIS.**

DISTINGUISHED EXPERT SERIES

- Mechanisms and predictors of high-risk human papillomavirus (HPV) clearance in the uterine cervix** 337
 K. Syrjänen - *Turku, FINLAND*
 This review discusses the mechanism and predictors of oncogenic human papillomavirus (HR-HPV) clearance in the uterine cervix.

ORIGINAL ARTICLES

- Radiation reduces carboplatin sensitivity and enhances nedaplatin sensitivity in cervical squamous cell carcinoma *in vitro*** 352
 T. Tanaka, K. Yukawa, N. Umesaki - *Wakayama, JAPAN*
 Radiation reduces carboplatin sensitivity but enhances nedaplatin sensitivity in cervical squamous cancer cells *in vitro*.
- Expression of matrix metalloproteinase-2 in serous borderline ovarian tumors is associated with non-invasive implant formation** 356
 T. Paulsen, A.H. Ree, J. Kærn, K. Kjærheim, A. Bassarova, A. Berner, T. Haldorsen, C. Tropé, J.M. Nesland - *Oslo, NORWAY*
 S-BOT with strong positive MMP-2 staining was frequently found in women with primary S-BOT and noninvasive implants.
- Feasibility and efficacy of chemotherapy with gemcitabine mono and with paclitaxel/mitoxantron in gynaecological cancers** 364
 C. Villena-Heinsen, A. Schroer, D. Lüdders, T. Cordes, N. Maass, M. Friedrich, D. Fischer - *Kiel, GERMANY*
 Gemcitabine mono in a 30 minute-schedule with the combination of paclitaxel/mitoxantron is a well tolerable regimen in advanced gynaecological cancers.
- Expression of the CXCR4 and CCR7 chemokine receptors in human endometrial cancer** 370
 J. Kodama, Hasengaowa, N. Seki, T. Kusumoto, Y. Hiramatsu - *Huhehaote, CHINA*
 CXCR4 and CCR7 are suppressed in high-grade endometrial cancers and may not play a crucial role in tumor progression.
- Comparison of symptoms and presentation of women with benign, low malignant potential and invasive ovarian tumors** 376
 C.M. Olsen, J. Cnossen, A.C Green, P.M. Webb - *Groningen, THE NETHERLANDS*
 A comparison of symptoms and presentation of women with benign, low malignant potential and invasive ovarian tumors is described.
- The role of three-dimensional (3D) sonography and 3D power Doppler in the preoperative assessment of borderline ovarian tumors** 381
 K. Kalmantis, Th. Papageorgiou, A. Rodolakis, E. Lymberopoulos, G. Daskalakis, Z. Voulgaris, A. Antsaklis - *Athens, GREECE*
 The value of 3D sonography and 3D power Doppler in distinguishing borderline ovarian tumors from benign cysts and malignant tumors was studied.

- Significance of hypoxia in uterine cervical cancer. Multicentre study** 386
 J. Markowska, J.P. Grabowski, K. Tomaszewska, Z. Kojs, J. Pudelek, M. Skrzypczak, J. Sobotkowski, J. Emerich, A. Olejek, V. Filas - *Poznan, POLAND*
 The adverse effects of radiotherapy according to expression levels of hypoxic markers in cervical cancer tissue were studied.
- Frequency of cervical intraepithelial neoplasia and infectious agents for vaginitis in menstrual cycle phase** 389
 R.S. Nomelini, P.L. Pansani, E.F.C. Murta - *Minas Gerais, BRAZIL*
 The frequency of cytolytic flora and *Candida albicans* can be influenced by the menstrual cycle phase, but CIN is not.
- Use of lysophosphatidic acid in the management of benign and malignant ovarian tumors** 394
 B. Požlep, M. Meleh, B. Kobal, I. Verdenik, J. Osredkar, L.Z. Kralj, H. Meden-Vrtovec - *Ljubljana, SLOVENIA*
 The value of lysophosphatidic acid determination in patients with benign and malignant ovarian tumors was examined.
- Hysteroscopy with directed biopsy versus dilatation and curettage for the diagnosis of endometrial hyperplasia and cancer in perimenopausal women** 400
 R. Bedner, I. Rzepka-Górska - *Szczecin, POLAND*
 Hysteroscopic-guided biopsy of the endometrium for histological examination is the most effective method for women at risk of endometrial cancer.
- CASE REPORTS**
- Ovarian clear cell carcinoma associated with endometriosis: a case report with immunohistochemical study** 403
 E.F.C. Murta, R.S. Nomelini, F.A. Ferreira, M.A. Lima - *Uberaba-MG, BRAZIL*
 The association of endometriosis and ovarian clear cell carcinoma should be kept in mind in patients with ovarian cysts.
- Acute abdomen in the postoperative period after cytoreductive surgery - case report** 406
 W.A.A. Tjalma, P. Elst, F. Ahankour - *Antwerp, BELGIUM*
 Acute abdomen in the postoperative period after cytoreductive surgery is a life threatening situation.
- Successful spontaneous pregnancy in a patient with rectal carcinoma treated with pelvic radiotherapy and concurrent chemotherapy: the unique role of laparoscopic lateral ovary transposition** 408
 M. Kurt, G. Uncu, S.K. Cetintas, N. Kucuk, S. Guler, L. Ozkan - *Bursa, TURKEY*
 Efficacy of laparoscopic lateral ovary transposition before pelvic radiotherapy in rectal cancer.
- Primary signet-ring cell carcinoma of the uterine cervix with long term follow-up: case report** 411
 L. Insabato, S. Simonetti, R. De Cecio, S. Di Tuoro, G. Bifulco, A. Di Spiezio Sardo - *Naples, ITALY*
 A case of primary signet-ring cell adenocarcinoma of the uterine cervix without evidence of recurrence eight years after the diagnosis is described.
- Salivary duct carcinoma presenting with vaginal metastasis: case report** 415
 M.Y. Gabriel, G.M. Yousef - *Toronto, CANADA*
 A case of a woman with salivary duct carcinoma presenting with vaginal metastasis is reported.
- A rare case of female pelvic mass: angioleiomyoma of the broad ligament** 418
 L. Cobellis, E. Pecori, F. Rigatti, C. Scaffa, M. Rotondi, E.M. Messalli - *Naples, ITALY*
 A rare case of angioleiomyoma of the broad ligament of the uterus is presented.

- A malignant eccrine poroma in a pregnant woman: case report and review of the literature** 421
E. Kairi-Vassilatou, D. Grapsa, N. Dafnios, V. Smyrniotis, D. Hasiakos, A. Kondi-Pafiti - *Athens, GREECE*
The first reported case of a malignant eccrine poroma in a pregnant woman is presented.
- Ovarian carcinoma as an incidental finding during cesarean section in a preeclamptic woman: case report** 423
Y. Engin-Ustun, Y. Ustun, K. Doğan, M.M. Meydanlı - *Malatya, TURKEY*
A rare case of advanced ovarian carcinoma diagnosed during cesarean section in a preeclamptic woman is presented.
- Liposarcoma of the breast - a case report** 425
B.C. Parikh, A. Ohri, M.Y. Desai, S.J. Pandya, R.I. Dave - *Gujarat, INDIA*
A case of liposarcoma of the breast is reported together with the different diagnostic and therapeutic modalities.

EUROPEAN JOURNAL OF GYNAECOLOGICAL ONCOLOGY

Vol. XXVIII, Number 5, 2007

ISSN: 0392-2936

I.R.O.G. CANADA, Inc.

Publisher's office: 4900 Côte St. Luc - Apt. 212 - Montréal, Québec H3W 2H3 (Canada) - Tél.: (514) 4893640-4893242 - Fax: (514) 4854513

E-mail: canlux@mgroup-online.com - www.irog.net

ALL RIGHTS RESERVED

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopies, recordings, nor any information storage and retrieval system without written permission from the copyright owner.

Printed in Italy