

CLINICAL AND EXPERIMENTAL OBSTETRICS & GYNECOLOGY

an International Journal

Founding Editor

A. Onnis

Montréal (CND)

Editors-in-Chief

M. Marchetti

Montréal (CND)

J.H. Check

Camden, NJ (USA)

Assistant Editor

J. Wilson

San Diego - CA (USA)

Editorial Board

Audet-Lapointe P., *Montréal (Canada)*

Axt-Fliedner R., *Lübeck (Germany)*

Basta A., *Krakow (Poland)*

Bender H.J., *Dusseldorf (Germany)*

Bhattacharya N., *Calcutta (India)*

Bonilla Musoles F., *Valencia (Spain)*

Charkviani T., *Tbilisi (Georgia)*

Dexeus S., *Barcelona (Spain)*

Di Paola G., *Buenos Aires (Argentina)*

Eskes T.K.A.B.,

Nijmegen (The Netherlands)

Franchi M., *Verona (Italy)*

Friedrich M., *Homburg (Germany)*

Gomel V., *Vancouver (Canada)*

Gorins A., *Paris (France)*

Grella P.V., *Padua (Italy)*

Holub Z., *Kladno (Czech Republic)*

Jordan J.A., *Birmingham, England (UK)*

Kaplan B., *Petach Tikva (Israel)*

Kralj B., *Ljubljana (Slovenia)*

Markowska J., *Poznan (Poland)*

Marth C., *Innsbruck (Austria)*

Meden-Vrtovec H., *Ljubljana (Slovenia)*

Ohara N., *Kobe (Japan)*

Papadopoulos N., *Alexandroupolis (Greece)*

Rakar S., *Ljubljana (Slovenia)*

Sciarra J.J., *Chicago, IL (USA)*

Stelmachow J., *Warsaw (Poland)*

Varras M.N., *Athens (Greece)*

Vîrtej P., *Bucharest (Romania)*

Winter R., *Graz (Austria)*

Publishing Organization (M. Morsani):

I.R.O.G. CANADA, Inc. - 4900 Côte St-Luc - Apt # 212 - Montréal, Qué. H3W 2H3 (Canada)

Tel. +514-4893242 - Fax +514-4854513 - E-mail: canlux@mgroun-online.com - www.irog.net

Editorial Office (M. Critelli):

Galleria Storione, 2/A - 35123 Padua (Italy) - Tel. (39) 049 8756900 - Fax (39) 049 8752018

CLINICAL AND EXPERIMENTAL OBSTETRICS AND GYNECOLOGY (ISSN 0390-6663) publishes original work, preferably brief reports, in the fields of Gynecology, Obstetrics, Fetal Medicine, Gynecological Endocrinology and related subjects. (Fertility and Sterility, Menopause, Uro-gynecology, Ultrasound in Obstetrics and Gynecology, Sexually Transmitted Diseases, Reproductive Biological Section). The Journal is covered by **INDEX MEDICUS, MEDLINE, EMBASE/Excerpta Medica**.

CLINICAL AND EXPERIMENTAL OBSTETRICS AND GYNECOLOGY is issued every three months in one volume per year by IROG CANADA Inc. Montréal. Printed in Italy by "La Garangola", Tipografia Editrice - Via E. Dalla Costa, 6 - 35129 Padova (Italy).

BRIEF COMUNICATION

Amici-Larciprete knotless technique. An easy way to suture in laparoscopy 69

G. Larciprete, F. Rossi, E. Valli, A. Amici, E. Cirese - *Rome, ITALY*

A new technique of suture in laparoscopy is described.

EDITORIAL ARTICLES

The concept and treatment methodology for inducing ovulation in women in apparent premature menopause 70

J.H. Check - *Camden, NJ (USA)*

By lowering elevated serum follicle stimulating hormone levels ovulation induction and pregnancies can be achieved despite the clinical appearance of menopause.

A putative role of versican in uterine leiomyomas 74

N. Ohara - *Kobe, JAPAN*

A putative role of versican in uterine leiomyomas is described.

ORIGINAL ARTICLES

Reproductive Biology Section

The effect of length of the follicular phase on pregnancy outcome following single embryo transfer (ET) in hypergonadotropic women 76

J.H. Check, B. Katsoff, D. Brasile, C. Wilson, J.K. Choe, J. Amui - *Camden, NJ (USA)*

In contrast to non-in vitro fertilization cycles a short follicular phase was not associated with a lower pregnancy rate in women treated with ethinyl estradiol.

High doses of GnRH antagonists are efficient in the management of severe ovarian hyperstimulation syndrome 78

F.M. Bonilla-Musoles, F. Raga, J.C. Castillo, M. Sanz, M. Dolz, N. Osborne - *Panama City, PANAMA*

Treatment with 3.0 mg of GnRH antagonists seems to be effective in the management of severe OHSS. Paracentesis was not required.

Repeated intracyclic clomiphene citrate therapy can be more effective than hMG therapy in inducing ovulation: case report 82

M. Kawamura, Y. Aoki, R. Kanda, H. Nakagomi, Y. Tsukahara, N. Hashimura - *Tokyo, JAPAN*

Repeated intracyclic clomiphene citrate therapy was more effective than high-dose hMG therapy in inducing ovulation in a patient with secondary amenorrhea.

No association found between decreased ovarian reserve and low thyroid function 85

J.H. Check, J. Stegonshek, C. Wilson - *Camden, NJ (USA)*

No association was found between diminished egg reserve and diminished thyroid function, suggesting that autoimmune ovarian damage is not the main common etiologic factor.

General Section

Factors affecting bone mineral density of young women and predictive factors of low bone mineral density 87

K. Ikeuchi, N. Umesaki - *Wakayama, JAPAN*

In young women BW and bone turnover markers were correlated with BMD. BW and NTx were predictive factors for low BMD.

Activity of telomerase in ovarian cancer cells. Clinical implications 91

J. Lubin, J. Markowska, A. Markowska, J. Stanisławiak, T. Lukaszewski - *Poznan, POLAND*

Telomerase activity in cell nuclei of ovarian cancer may be an independent prognostic index.

Comparison of the number of uterine myomas detected by in-office transvaginal ultrasonography removed by laparotomic myomectomy: preoperative work-up concerns	97
M. Fambrini, F. Tondi, G. Scarselli, C. Penna, A. Pieralli, K.L. Andersson, M. Marchionni - <i>Florence, ITALY</i> In-office TVS overlooks at least one myoma in one-third of patients theoretically eligible for minimally invasive conservative surgery.	
Catalase activity, serum trace element and heavy metal concentrations, vitamin A, vitamin D and vitamin E levels in hydatidiform mole	102
A. Kulusari, E. Adali, M. Kurdoglu, R. Yildizhan, A. Cebi, T. Edirne, H. Demir, I.H. Yoruk - <i>Van, TURKEY</i> Serum levels of catalase, Zn, Co, and vitamin A, D, and E were significantly lower in patients with hydatiform mole compared with healthy pregnant women and non pregnant controls.	
The contribution of anaesthesia modus on reducing blood loss during caesarean section	105
V. Liberis, P. Tsikouras, T. Vogiatzaki, A. Savidis, A. Ammari, Ch. Iatrou, G. Maroulis - <i>Thrace, GREECE</i> The trend from general anaesthesia to regional anaesthesia (RA) for caesarean section continues and RA is associated with lower levels of blood loss.	
Lipid peroxidation and antioxidant activity in complicated pregnancies	110
S.B. Patil, M.V. Kodliwadmath, S.M. Kodliwadmath - <i>Karnataka, INDIA</i> Oxidative stress in preeclampsia and eclampsia leads to decreased activity of antioxidants.	
The presence of <i>Helicobacter pylori</i> in cervical preinvasive lesions	113
T. Kaya, H. Ozan, C. Özakin - <i>Bursa, TURKEY</i> The cervix was investigated for the presence of <i>Helicobacter pylori</i> and found free of bacteria even in patients with active infection.	
Fetuses with single umbilical artery: analysis of 45 cases	116
B. Dane, C. Dane, M. Kiray, A. Cetin, M. Yayla - <i>Istanbul, TURKEY</i> Scanning the umbilical cord is one of the essential parts of ultrasonographic examination. The newborn should be reexamined immediately after birth due to the possibility of associated-undetected anomalies.	
Is pregnancy over 45 with very high parity related with adverse maternal and fetal outcomes?	120
A. Kale, U. Kuyumcuoğlu, A.Güzel - <i>Diyarbakir, TURKEY</i> Older women with very high parity were studied in relation to adverse maternal and fetal outcomes.	
CASE REPORTS	
Successful long-term management of adenomyosis associated with deep thrombosis by low-dose gonadotropin-releasing hormone agonist therapy	123
S. Akira, N. Iwasaki, M. Ichikawa, K. Mine, Y. Kuwabara, T. Takeshita, H. Tajima - <i>Tokyo, JAPAN</i> Low-dose GnRH _a therapy can maintain therapeutic effectiveness against adenomyosis for a long period, while suppressing the expression of adverse events.	
Conservative treatment by endoscopy of a cesarean scar pregnancy: two case reports	126
C. Colomé, M.T. Cusidó, L. Hereter, M.A. Pascual, R. Fábregas - <i>Barcelona, SPAIN</i> Our experience in the management of cesarean scar pregnancy with endoscopic surgery of two patients who desired to preserve their fertility is presented.	
Ruptured cornual pregnancy: case report	130
T. Takei, S. Matsuoka, N. Ashitani, N. Makihara, M. Morizane, N. Ohara - <i>Kobe, JAPAN</i> An uncommon case of ruptured cornual pregnancy occurring at 12 weeks of gestation is reported.	
A case report supporting the concept that some women have a predisposition for maternal meiosis errors resulting in digyny	133
J.H. Check, B. Katsoff, D. Summers-Chase, J. Breitbart - <i>Camden, NJ (USA)</i> Evaluation of a woman with two documented miscarriages related to triploidy by in vitro fertilization and pre-implantation genetic diagnosis showed that some women may have a predisposition for failure of polar body extrusion.	
Mucocele of the appendix mimicking an adnexal mass: a case report	135
D. Vavilis, D. Athanatos, D. Tsolakidis, A. Papanicolaou, J.N. Bontis - <i>Thessaloniki, GREECE</i> Appendiceal mucocele mimicking adnexal tumour.	