

europaean journal of gynaecological oncology

an International Journal

Editors-in-Chief

A. Onnis **P. Bószé**
Montréal (Canada) *Budapest (Hungary)*

Associate Editor

M. Marchetti
Padua (Italy)

Assistant Editor

J. Wilson
San Diego - CA (USA)

Editorial Board

H. H. Allen, <i>London, Ontario (Canada)</i>	A. F. Fuller Jr., <i>Boston, MA (USA)</i>	Leon F. A. G. Massuger, <i>Nijmegen (The Netherlands)</i>
S. O. Anteby, <i>Jerusalem (Israel)</i>	H. E. Geisler, <i>Indianapolis, IN (USA)</i>	J. Menczer, <i>Savyon (Israel)</i>
P. Audet-Lapointe, <i>Montreal (Canada)</i>	A. Gerbaulet, <i>Villejuif (France)</i>	J. Monsonego, <i>Paris (France)</i>
L. Aurelian, <i>Baltimore, MD (USA)</i>	A. Gorins, <i>Paris (France)</i>	A. Pinotti, <i>São Paulo (Brasil)</i>
A. Ayhan, <i>Ankara (Turkey)</i>	A.P.M. Heintz, <i>Utrecht (The Netherlands)</i>	S. M. Piver, <i>Buffalo, NY (USA)</i>
O. Balat, <i>Graziantep (Turkey)</i>	L. Ioannidou-Mouzaka, <i>Athens (Greece)</i>	S. Rakar, <i>Ljubljana (Slovenia)</i>
H. K. Barber, <i>New York, NY (USA)</i>	J. A. Jordan, <i>Birmingham, England (UK)</i>	K. S. Raju, <i>London, England (UK)</i>
A. Basta, <i>Krakow (Poland)</i>	A. I. Karseladze, <i>Moscow (Russia)</i>	J. H. Shepherd, <i>London, England (UK)</i>
H. C. Bender, <i>Dusseldorf (Germany)</i>	J. Klustersky, <i>Bruxelles (Belgium)</i>	P. Sismondi, <i>Turin (Italy)</i>
P. L. Benedetti Panici, <i>Rome (Italy)</i>	V. Kozachenko, <i>Moscow (Russia)</i>	J. Stelmachów, <i>Warsaw (Poland)</i>
T. Charkviani, <i>Tbilisi (Georgia)</i>	B. Kralj, <i>Ljubljana (Slovenia)</i>	L. Ungár, <i>Budapest (Hungary)</i>
D. Dargent, <i>Lyon (France)</i>	E. Kubista, <i>Vienna (Austria)</i>	I. Vergote, <i>Leuven (Belgium)</i>
C. F. De Oliveira, <i>Coimbra (Portugal)</i>	A. P. Kudelka, <i>Houston, TX (USA)</i>	J. B. Vermorken, <i>Edegem (Belgium)</i>
S. Dexeus Jr., <i>Barcelona (Spain)</i>	T. Maggino, <i>Padua (Italy)</i>	P. Vírtej, <i>Bucharest (Romania)</i>
G. De Palo, <i>Milan (Italy)</i>	J. Markowska, <i>Poznan (Poland)</i>	P.-H. Wang, <i>Taipei (Taiwan)</i>
G. R. Di Paola, <i>Buenos Aires (Argentina)</i>	C. Marth, <i>Innsbruck (Austria)</i>	R. Winter, <i>Graz (Austria)</i>
F. Di Re, <i>Milan (Italy)</i>		Y. Yokoyama, <i>Hirosaki (Japan)</i>
P. Di Saia, <i>Orange, CA (USA)</i>		J. Zielinski, <i>Warsaw (Poland)</i>
S. K. Fotiou, <i>Athens (Greece)</i>		

Publisher Organization:

I.R.O.G. CANADA, Inc. - 4900 Côte St-Luc
Apt # 212 - Montréal, Qué. H3W 2H3 (Canada)
Tel. +514-4893242-4893640 - Fax +514-4854513 - E-mail: canlux@qc.aira.com
www.irog.net

EUROPEAN JOURNAL OF GYNAECOLOGICAL ONCOLOGY (ISSN 0392-2936) publishes original peer reviewed works in the fields of female genital cancers and related subjects and also proceedings of gynecologic oncology society meetings all over the world. The Journal is covered by **CURRENT CONTENTS, SCISEARCH, RESEARCH ALERT, INDEX MEDICUS, MEDLINE, EMBASE/Excerpta Medica, CURRENT ADVANCES IN CANCER RESEARCH, BIOSIS.**

DISTINGUISHED EXPERT SERIES

Progestins and risk of breast cancer. Much ado about nothing? 345

A. Gorins - *Paris, FRANCE*

A very small increased risk of breast cancer with estrogen therapy has been shown; however bias are often present. It has not been clearly demonstrated that the adjunction of progestin to estrogen significantly increases the risk, nor that it brings about a protective effect. Some recent works seem to indicate that continuous estroprogestin replacement therapy might be deleterious, while a sequential regimen would not be. This is to be confirmed. The data and conclusions of the medical literature should be interpreted very cautiously.

The current status of HPV DNA testing 351

N. J. Agnantis, A. Sotiriadis, E. Paraskevaidis - *Ioannina, GREECE*

HPV testing may be useful in certain cases of screening, in a triage of women with ASCUS and in post-treatment surveillance of women with CIN.

EDITORIAL ARTICLE

How to avoid the uncertainties of intraoperative examination of the sentinel lymph node in breast cancer? 357

Y. Aubard, J. Mollard, V. Fermeaux - *Limoges, FRANCE*

A new strategy for the detection of a sentinel lymph node (SLN) under local anaesthesia (SLNLA) before tumorectomy is proposed. This avoids any second operation for axillary lymphadenectomy and a wrongful lymphadenectomy in cases of false positives.

ORIGINAL ARTICLES

Methylation: A second hit in the two-hit hypothesis 361

J. P. Geisler, J. A. Rathe, K. J. Manahan, H. E. Geisler - *Indianapolis, IN (USA)*

Methylation of one BRCA1 allele may serve as the second hit in a patient with a diploid locus and missense mutation on the other allele.

Cancer of the cervix with endometrial extension treated by radiotherapy: A retrospective case-control study 362

F. L. Ampil, G. Caldito, J. B. Unger - *Louisiana, USA*

The prognosis of cancer of the cervix is linked to tumor stage. A case-control retrospective study showed a worse prognosis in patients with endometrial extension of cervical cancer.

Development of an evidence-based algorithm for the management of cervical cancer 365

M. C. Shaw, C. D. A. Wolfe, O. Devaja, K. S. Raju - *London, UK*

The development of an algorithm for cervical cancer management and the level of supporting primary research evidence are described.

High-risk human papillomavirus type does not predict grade of cervical intraepithelial neoplasia 373

M. F. Evans, S. L. Mount, P. M. Vacek, K. Cooper - *Burlington, VT (USA)*

Predicting CIN grade on the basis of HPV testing following smear diagnosis may be improved by redefining the "high-risk" HPV type classification.

Successful treatment with weekly paclitaxel in a patient with recurrent endometrial cancer. A case report 377

S-J Chang, T-C Chen, T-Y Wang, C-Y Hsu, H-S Chen, Y-C Yang - *Taipei, TAIWAN*

A case of recurrent endometrial cancer is reported. The patient responded well to weekly paclitaxel, as demonstrated by serial power Doppler angiography.

Laparoscopic gonadectomy in a case of a dicentric fluorescent Y-chromosome mosaicism with Turner-like phenotype and virilized external genitalia 379

K. Takeuchi, S. Oomori, N. Oda, S. Takekida, H. Kondo, T. Maruo - *Kakogawa, JAPAN*

Laparoscopic gonadectomy was performed in a case of a dicentric fluorescent Y-chromosome mosaicism with Turner-like phenotype and virilized external genitalia.

Endometrial cancer: factors affecting survival	381
R. Labastida, S. Dexeus, R. Fábregas, F. Tresserra, A. Fernández - <i>Barcelona, SPAIN</i>	
Lack of myometrial invasion, negative lymph nodes and age under 60 are significant predicting factors of survival in endometrial cancer.	
Primary squamous cell carcinoma of the breast	384
K. Brack, P. C. G. I. Ramaekers, W. A. A. Tjalma - <i>Edegem, BELGIUM</i>	
Primary squamous cell carcinoma of the breast is a rare variant of a metaplastic carcinoma. The treatment approach of this rare disease is discussed.	
Identification of the sentinel lymph node stained with blue dye in breast cancer patients	387
A. C. Rufino, F. F. Baracat, A. P. Madeiro, U. G. Lippi - <i>São Paulo, BRAZIL</i>	
The use of blue dye in sentinel lymphadenectomy of breast cancer cases offers adequate identification of sentinel lymph nodes, though it does not determine accurate staging.	
Laparoscopically assisted modified radical vaginal hysterectomy (type II) with pelvic lymphadenectomy: ultrasonic operative technique	391
Z. Holub, J. Lukáč, L. Kliment, Š. Urbánek - <i>Kladno, CZECH REPUBLIC</i>	
In five cases the surgical treatment of early cervical cancer was successfully performed using an ultrasonic operative technique during the laparoscopic phase of laparoscopically assisted radical vaginal hysterectomy type II and pelvic lymphadenectomy.	
Laparoscopic surgery in treatment of Stage IIb cervical cancer after neoadjuvant chemotherapy. A case report and review of the literature	393
M. Malzoni, V. Spina, G. Perniola, V. Aleandri, B. Mossa, T. Iuele, F. Imperato - <i>Rome, ITALY</i>	
Laparoscopic radical hysterectomy potentially allows for decreased perioperative morbidity and blood loss, faster recovery and better cosmetic results. Large studies with long-term follow-up are needed to confirm that this approach may be proposed as an alternative to conventional surgery.	
Pelvic and paraaortic lymph node metastasis in clinical Stage I endometrial adenocarcinoma: an analysis of 58 consecutive cases	398
M. Özsoy, S. Dilek, D. Ozsoy - <i>Isparta, TURKEY</i>	
A complete lymphadenectomy was done in 58 consecutive cases of clinical Stage I endometrial adenocarcinomas in order to determine the real stages of the cases.	
Pelvic inflammatory disease is a risk factor for cervical cancer	401
J. Skapinyecz, I. Smid, A. Horváth, Cs. Jeney, L. Kardos, P. Kovács - <i>Budapest and Debrecen, HUNGARY</i>	
Higher prevalence of human papillomavirus in patients with pelvic inflammatory disease suggests that this disease means a risk for cervical cancer.	
Is pretreatment hemoglobin level a predictor of complete response to salvage chemotherapy for recurrent platinum-pretreated ovarian carcinoma?	405
A. Gadducci, S. Cosio, A. Fanucchi, L. Tanganelli, P. F. Conte, R. Cristofani, A. R. Genazzani - <i>Pisa, ITALY</i>	
Baseline hemoglobin level is an independent predictor of complete response to salvage chemotherapy in patients with recurrent platinum-pretreated ovarian carcinoma.	
The relationship between expression of p53/bcl-2 and clinicopathological criteria in cervix squamous cell carcinomas	411
M. Bitiren, E. A. Cakmak, A. Gocmen, S. S. Inaloz, I. Sari, M. Karakok, A. Aydin - <i>Gaziantep, TURKEY</i>	
A decrease in bcl-2 reactivity and an increase in p53 positive staining were noted as the histological differentiation worsened.	
The importance of fine needle aspiration biopsy and sonographic evaluation of parametria in cervical cancer	413
K. Cendrowski, W. Sawicki, B. Śpiewankiewicz, J. Stelmachów - <i>Warsaw, POLAND</i>	
Fine needle aspiration biopsy (FNAB) and transvaginal ultrasonography (TVS) are very useful methods in preoperative assessment of the parametria in cervical cancer. Correct staging qualification, especially evaluation of the parametria is very important in choosing of adequate method of treatment.	
Prognostic value of matrix metalloproteinase-9 (gelatinase-B) expression in epithelial ovarian tumors	417
S. Ozalp, H. M. Tanir, O. T. Yalcin, S. Kabukcuoglu, U. Oner, M. Uray - <i>Eskisehir, TURKEY</i>	
Stromal but not epithelial matrix metalloproteinase (MMP-9) expression predicts poor survival in epithelial ovarian cancer.	
Adjuvant chemotherapy following surgery in the management of uterine sarcomas	421
A. Szánthó, J. Bálega, I. Szabó, A. Demeter, N. Sipos, Z. Csapó, Z. Papp - <i>Budapest, HUNGARY</i>	
Adjuvant chemotherapy according to the CYVADIC-protocol for uterine sarcoma patients gave an overall median time to progression of 22.14 months and good results for the patients with lung metastases.	

Microvessel density as a prognostic factor in preinvasive and invasive cervical lesions	425
S. Ozalp, O. T. Yalcin, U. Oner, H. M. Tanir, M. Acikalın, I. Sarac - <i>Eskisehir, TURKEY</i>	
Microvessel density, a marker for angiogenesis that is involved in the development of cervical neoplasms, may be used as a prognostic factor.	
Ovarian fibroma: our experience of 34 cases	429
G. Gargano, M. De Lena, F. Zito, G. Fanizza, V. Mattioli, F. Schittulli - <i>Bari, ITALY</i>	
This retrospective study examines the percentage of ovarian fibromas found in the Division of Pathology Oncology Institute of Bari from January 1994 to December 2001.	
Failure of sentinel node identification following neo-adjuvant chemo-radiotherapy for locally advanced squamous cell carcinoma of the vulva	433
H. Levavi, G. Sabah, R. Hardoff, C. Koren, H. Gutman - <i>Petah Tiqva and Ramat Aviv, ISRAEL</i>	
Lymphoscintigraphy + dye - sentinel node detection in neo-adjuvant chemo-radiotherapy vulvar cancer failed due to irradiation damage to lymph channels.	
CASE REPORTS	
A case of adenocarcinoma of the endometrial type mixed with a clear cell component that metastasized to the vagina	435
Y. Yokoyama, A. Sakamoto, M. Umemoto, M. Futagami, T. Sakamoto, H. Maruyama, S. Sato, H. Mizunuma - <i>Hirosaki, JAPAN</i>	
A rare case of endometrial-type adenocarcinoma mixed with a clear cell component that metastasized to the vagina is presented.	
Clear cell ovarian carcinoma in a pregnant woman with a history of infertility, endometriosis and unsuccessful IVF treatment	438
G. Makrydimas, A. Sotiriadis, E. Paraskevaidis, N. Pavlidis, N. Agnantis, D. Lolis - <i>Ioannina, GREECE</i>	
A case of a 37-year-old pregnant woman with clear cell ovarian carcinoma and a history of infertility, endometriosis and IVF is presented.	
Sclerosing stromal tumor of the ovary: A case report	442
E. Kuscu, M. Oktem, H. Karahan, B. Bilezikci, B. Demirhan - <i>Ankara, TURKEY</i>	
A 34-year-old woman presented with hirsutism, oligomenorrhea of three months' duration, and a palpable right adnexal mass. At surgery the mass was found to originate from the left ovary. Left oophorectomy was performed, and the lesion was identified as a sclerosing stromal tumor of the ovary.	
A ruptured ovarian endometrioma mimicking ovarian malignancy: case report	445
A. Göçmen, M. Karaca, M. Tarakçioğlu - <i>Gaziantep, TURKEY</i>	
A case of ruptured ovarian endometrioma thought to be a malignant ovarian tumor because of extremely high CA 125 levels is presented.	
Clitoromegaly in type 2 neurofibromatosis: A case report and review of the literature	447
H. Yüksel, A. R. Odabaşı, S. Kafkas, E. Onur, M. Turgut - <i>Aydın, TURKEY</i>	
To our knowledge this is the first case report of clitoromegaly with type 2 neurofibromatosis in the English literature.	
Books Review	452