

# europaean journal of gynaecological oncology

an International Journal

*Founding Editor*

**A. Onnis**

*Montréal (Canada)*

*Editors-in-Chief*

**M. Marchetti**

*Montréal (Canada)*

**P. Bószé**

*Budapest (Hungary)*

*Associate Editor*

**T. Maggino**

*Padua (Italy)*

*Assistant Editor*

**J. Wilson**

*San Diego - CA (USA)*

*Editorial Board*

- |   |  |   |
|---|--|---|
| Allen H.H., <i>London, Ontario (Canada)</i>       | Geisler H.E., <i>Indianapolis, IN (USA)</i>                | Monsonog J., <i>Paris (France)</i>            |
| Ayhan A., <i>Ankara (Turkey)</i>                  | Gorins A., <i>Paris (France)</i>                           | Pálfalvi L., <i>Budapest, (Hungary)</i>       |
| Balat O., <i>Graziantep (Turkey)</i>              | Heintz A.P.M.,<br><i>Utrecht (The Netherlands)</i>         | Piura B., <i>Beer Sheva (Israel)</i>          |
| Bănceanu G., <i>Bucarest (Romania)</i>            | Ioannidou-Mouzaka L.,<br><i>Athens (Greece)</i>            | Piver S.M., <i>Buffalo, NY (USA)</i>          |
| Basta A., <i>Krakow (Poland)</i>                  | Jordan J.A.,<br><i>Birmingham, England (UK)</i>            | Rakar S., <i>Ljubljana (Slovenia)</i>         |
| Bender H.C., <i>Dusseldorf (Germany)</i>          | Klastersky J., <i>Bruxelles (Belgium)</i>                  | Shepherd J.H.,<br><i>London, England (UK)</i> |
| Charkviani T., <i>Tbilisi (Georgia)</i>           | Kubista E., <i>Vienna (Austria)</i>                        | Smit B.J., <i>Tygerberg (South Africa)</i>    |
| Chiarelli S., <i>Padua (Italy)</i>                | Lee Y.S., <i>Daegu (South Korea)</i>                       | Stelmachów J., <i>Warsaw (Poland)</i>         |
| De Oliveira C.F., <i>Coimbra (Portugal)</i>       | Markowska J., <i>Poznan (Poland)</i>                       | Syrjänen K., <i>Turku (Finland)</i>           |
| Dexeus S. Jr., <i>Barcelona (Spain)</i>           | Marth C., <i>Innsbruck (Austria)</i>                       | Tjalma W., <i>Antwerpen (Belgium)</i>         |
| Di Paola G.R.,<br><i>Buenos Aires (Argentina)</i> | Massuger Leon F.A.G.,<br><i>Nijmegen (The Netherlands)</i> | Ungár L., <i>Budapest (Hungary)</i>           |
| Di Re F., <i>Milan (Italy)</i>                    | Menczer J., <i>Savyon (Israel)</i>                         | Vermorken J.B., <i>Edegem (Belgium)</i>       |
| Di Saia P., <i>Orange, CA (USA)</i>               |  | Wang P.H., <i>Taipei (Taiwan)</i>             |
| Elit L., <i>Hamilton (Canada)</i>                 |  | Winter R., <i>Graz (Austria)</i>              |
| Friedrich M., <i>Hamburg (Germany)</i>            |  | Yokoyama Y., <i>Hirosaki (Japan)</i>          |

*Publishing Organization (M. Morsani):*

I.R.O.G. CANADA, Inc. - 4900 Côte St-Luc - Apt # 212 - Montréal, Qué. H3W 2H3 (Canada)  
Tel. +514-4893242 - Fax +514-4854513 - E-mail: canlux@mgroup-online.com - www.irog.net

*Editorial Office (M. Critelli):*

Galleria Storione, 2/A - 35123 Padua (Italy) - Tel. (39) 049 8756900 - Fax (39) 049 8752018

EUROPEAN JOURNAL OF GYNAECOLOGICAL ONCOLOGY (ISSN 0392-2936) publishes original peer reviewed works in the fields of female genital cancers and related subjects and also proceedings of gynecologic oncology society meetings all over the world. The Journal is covered by **CURRENT CONTENTS, SCISEARCH, RESEARCH ALERT, INDEX MEDICUS, MEDLINE, EMBASE/Excerpta Medica, CURRENT ADVANCES IN CANCER RESEARCH, BIOSIS.**



# Contents

European Journal of Gynaecological Oncology - Vol. XXXI, no. 3, 2010

## DISTINGUISHED EXPERT SERIES

- Could the eventual results of the NSABP\* 39/RTOG\*\* 0413 trial for partial breast irradiation (PBI) be improved by combining spherical applicators and whole breast irradiation? Radiobiology suggests it may** 241
- B.J. Smit - *Tygerber, SOUTH AFRICA*  
Combining spherical application and whole breast irradiation may improve outcome of partial breast irradiation.

## EXPERT'S POINT OF VIEW

- Management of abnormal cytological findings** 250
- P. Bösze - *Budapest, HUNGARY*  
Point of view of experts on management of abnormal cytological findings is discussed.

## REVIEW ARTICLES

- Hyperthermic intraperitoneal chemotherapy added to the treatment of ovarian cancer. A review of achieved results and complications** 256
- E. Dovern, I.H.J.T. de Hingh, V.J. Verwaal, W.J. van Driel, S.W. Nienhuijs - *Eindhoven, THE NETHERLANDS*  
A review of achieved results and complications of hyperthermic intraperitoneal chemotherapy added to the treatment of ovarian cancer.
- Hormone therapy/adjuvant chemotherapy induced deleterious effects on the bone mass of breast cancer patients and the intervention of physiotherapy: a literature review** 262
- T. Tonzzer, C.M.A.P. Pereira, U.P. Filho, A. Marx - *São Paulo, BRAZIL*  
The aim of this study was a literature review. Twenty four articles were selected to analyze the possible deleterious effects caused by adjuvant chemotherapy and hormone therapy on the bone mass of patients diagnosed with breast cancer and the physiotherapy intervention.
- Molecular markers in epithelial ovarian cancer: Their role in prognosis and therapy** 268
- F. Zagouri, M.A. Dimopoulos, E. Bournakis, C.A. Papadimitriou - *Athens, GREECE*  
The use of molecular markers in epithelial ovarian cancer in common clinical practice seems promising for the diagnosis and prognostication. It is tempting to anticipate the gradual integration of molecular profiling in clinical practice.

## ORIGINAL ARTICLES

- Human papillomavirus (HPV)-type distribution in relation to oral contraceptive use in women with cervical intraepithelial neoplasia, Durban, South Africa** 278
- M. Moodley, G. Lindeque, C. Connolly - *Durban, SOUTH AFRICA*  
Unusual HPV types with contraceptive usage in a South African context.
- A comparison of outcome in patients with Stage 1 clear cell and grade 3 endometrioid adenocarcinoma of the endometrium with and without adjuvant therapy** 284
- J.A. Rauh-Hain, I. Costaaggini, A.B. Olawaiye, W.B. Growdon, N.S. Horowitz, M.G. del Carmen - *Pittsburgh, USA*  
In surgical Stage I uterine clear cell carcinoma there is no clear benefit to adjuvant radiation given the absence of improvement in recurrence risk or any survival benefit.
- Liquid based cytology improves the positive predictive value of glandular smears compared to conventional cytology** 288
- R.J. Edmondson, C.A. Errington, D.J.A. Mansour - *Newcastle upon Tyne, UK*  
A database review suggests that LBC increases the positive predictive value of cervical sampling to detect preinvasive and invasive cancer.





<i>Contents</i>	239
<b>Increase of Mcm3 and Mcm4 expression in cervical squamous cell carcinomas</b>	291
N. Gan, Y. Du, W. Zhang, J. Zhou - <i>Zhejiang Province, CHINA</i> Mcm3 and Mcm4 were highly expressed in cervical squamous carcinoma and these two proteins might be useful as biomarkers in clinical diagnosis.	
<b>Repeated chemosensitivity testing in patients with epithelial ovarian carcinoma</b>	295
M. Pospíšková, M. Špenerová, R. Pilka, M. Kudela, M. Hajdúch, V. Šrámek, B. Melichar, K. Cwiertka - <i>Olomouc, CZECH REPUBLIC</i> Changes of chemosensitivity were repeatedly evaluated in a cohort of patients with epithelial ovarian carcinoma. Limited, but statistical increase in chemoresistance was observed for paclitaxel and carboplatin.	
<b>Non-hormonal treatment of vasomotor symptoms in gynecological cancer patients</b>	299
L. Del Pup, T. Maggino - <i>Mestre-Venice, ITALY</i> Non-hormonal therapies such as venlafaxine, paroxetine, gabapentin and clonidine have been shown to reduce hot flashes in randomized trials compared to placebo.	
<b>Analysis of clinical and molecular associations of triple negative breast cancers in node-negative patients</b>	304
N. Koronakis, G. Karanikas, E.E. Lagoudianakis, D. Grosomanidis, A. Pappas, J. Chrysikos, G. Xepapadakis, A. Manouras - <i>Athens, GREECE</i> Immunohistochemical expression of patients with node-negative invasive breast carcinomas was compared. The TN phenotype is associated with aggressive histology.	
<b>Bevacizumab, paclitaxel and carboplatin for advanced ovarian cancer: low risk of gastrointestinal and cardiovascular toxicity</b>	308
L.N. Abaid, K.L. Lopez, J.P. Micha, M.A. Rettenmaier, J.V. Brown III, B.H. Goldstein - <i>Newport Beach, CA (USA)</i> Despite concerns for hypertension and bowel perforations, bevacizumab can be safely administered during induction therapy for ovarian carcinoma.	
<b>Expression of p16 in serous ovarian neoplasms</b>	312
H.O. Nazlioglu, İ. Ercan, T. Bilgin, S. Ozuysal - <i>Bursa, TURKEY</i> P16 expression of benign, borderline and malignant serous ovarian neoplasms was immunohistochemically evaluated, and was found to be strong in serous ovarian carcinomas.	
<b>Distribution of HPV genotypes in uterine cervical lesions among the Uighur women in Xinjiang province of China</b>	315
A. Abudukadeer, Y. Ding, M. Niyazi, A. Ababaikeli, A. Abudula - <i>Urumqi, P.R. CHINA</i> HPV16 was the most common type found in Uighur patients with cervical cancer while HPV18 and 58 types were relatively low and other types were absent.	
<b>Intestinal-type metaplasia in the original squamous epithelium of the cervix</b>	319
E. Sivridis, G. Karpathiou, V. Malamou-Mitsi, A. Giatromanolaki - <i>Ioannina, GREECE</i> The expression of mucin-distended goblet cells in the normal/original squamous epithelium of the cervix is described for the first time.	
<b>Evaluation of preoperative diagnosis with results of histopathological examinations of ovarian tumors in women of reproductive age</b>	323
D. Samulak, S. Sajdak, M. Wilczak, M. Michalska, B. Pi ta, M. Englert-Golon - <i>Poznan, POLAND</i> Gynecological, ultrasonographic and Doppler examinations as well as determination of CA-125 antigen provide an efficient research panel in the preoperative diagnosis of ovarian tumors.	
<b>CASE REPORTS</b>	
<b>Management of recurrence from a retroperitoneal xanthogranuloma: case report</b>	326
D. Salehin, C. Haugk, A. Stricker, R. Triefenbach, M. Friedrich - <i>Krefeld, GERMANY</i> A rare case of a xanthogranuloma from the retroperitoneal space diagnosed in a 24-year-old woman is presented.	





- A benign metastasizing leiomyoma involving a nodule in the pulmonary artery: case and literature review** 329  
O. Poujade, A.S. Genin, M. Dhouha, D. Luton - *Paris, FRANCE*  
A benign metastasizing leiomyoma with local recurrence and metastasis growing into the left pulmonary artery was managed by surgery.
- Metastatic and recurrent adenocarcinoma of the uterine cervix: a long-term survival of 16 years** 333  
I. Peregrin-Alvarez, M.N. Akl, C.P. Morrow, J.F. Magrina - *Seville, SPAIN*  
A multi-modal salvage approach may achieve long-term survival in rare cases of recurrent metastatic cervical adenocarcinoma.
- Three-dimensional power Doppler color ultrasonographic features of a minimal deviation adenocarcinoma of the uterine cervix** 336  
L. Hereter, F. Tresserra, B. Graupera, M.A. Pascual, M.A. Martinez, A. Úbeda - *Barcelona, SPAIN*  
A case of minimal deviation adenocarcinoma of the cervix in a 34-year-old female is presented.
- Ovarian metastasis of a primary renal cell carcinoma: case report and review of literature** 339  
S. Guney, N. Guney, D. Özcan, T. Sayılğan, E. Özakın - *Istanbul, TURKEY*  
A case of ovarian metastasis of a primary renal cell carcinoma in a 29-year-old patient is reported.
- Liver resection for metastases arising from recurrent granulosa cell tumour of the ovary - a case series** 342  
T.K. Madhuri, S. Butler-Manuel, N. Karanjia, A. Tailor - *Guildford, UK*  
This series discusses three recurrent cases of GCT of the ovary successfully managed with resection of hepatic metastases improving patient quality of life.
- Ovarian carcinomatosis presenting as bilateral inguinal hernia: a brief report** 345  
Y.H. Hung, C.T. Hsu, C.C. Chang - *Taiwan, R.O.C.*  
Gynecologists should be aware of the difficulties associated with a delay in diagnosis of ovarian carcinomatosis presenting as inguinal hernias.
- Serous ovarian cystadenocarcinoma incidentally discovered in a 29-year-old patient: case report** 347  
N. Cutura, V. Soldo, M. Vasiljevic - *Belgrade, SERBIA*  
A case report of a serous ovarian cystadenocarcinoma incidentally discovered in a 29-year-old women is presented.
- Chemotherapy with low-dose bevacizumab and carboplatin in the treatment of a patient with recurrent cervical cancer** 350  
S.J. Tan, Y.H. Juan, P.T. Fu, M.H. Yu, H.C. Lai - *Taipei, TAIWAN*  
The case of patient with platinum-resistant recurrent cervical cancer treated with low-dose bevacizumab and carboplatin with resultant improved disease progression is presented.
- Groin recurrence following Stage IA squamous cell carcinoma of the vulva with negative nodes on superficial inguinal lymphadenectomy** 354  
A.C. Iyibozkurt, O.C. Dural, S. Topuz, S. Berkman, E. Bengisu - *Istanbul, TURKEY*  
Stage IA vulvar cancer may recur in the groin despite finding negative nodes in superficial inguinal lymph node dissection.

## EUROPEAN JOURNAL OF GYNAECOLOGICAL ONCOLOGY

Vol. XXXI, Number 3, 2010

ISSN: 0392-2936

I.R.O.G. CANADA, Inc.

Publisher's office: 4900 Côte St. Luc - Apt. 212 - Montréal, Québec H3W 2H3 (Canada) - Tél.: (514) 4893640-4893242 - Fax: (514) 4854513

E-mail: canlux@mgroup-online.com - www.irog.net

ALL RIGHTS RESERVED

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopies, recordings, nor any information storage and retrieval system without written permission from the copyright owner.

Printed in Italy

